

ON KREBES' TANGLE

SUSAN M. ABERNATHY

ABSTRACT. A genus-1 tangle $\mathcal{G}$ is an arc properly embedded in a standardly embedded solid torus S in the 3-sphere. We say that a genus-1 tangle embeds in a knot $K \subseteq S^3$ if the tangle can be completed by adding an arc exterior to the solid torus to form the knot K . We call K a closure of $\mathcal{G}$. An obstruction to embedding a genus-1 tangle $\mathcal{G}$ in a knot is given by torsion in the homology of branched covers of S branched over $\mathcal{G}$. We examine a particular example $\mathcal{A}$ of a genus-1 tangle, given by Krebs, and consider its two double-branched covers. Using this homological obstruction, we show that any closure of $\mathcal{A}$ obtained via an arc which passes through the hole of S an odd number of times must have determinant divisible by three. A resulting corollary is that if $\mathcal{A}$ embeds in the unknot, then the arc which completes $\mathcal{A}$ to the unknot must pass through the hole of S an even number of times.

1. INTRODUCTION

We choose a standardly embedded solid torus $S^1 \times D^2 \subset S^3$, denoted by S . Then a genus-1 tangle is a properly embedded arc in S . Just as we may discuss embedding ordinary tangles in B^3 into knots and links (see [1], [2], and [4]), we may consider embedding genus-1 tangles in knots. We say that a genus-1 tangle $\mathcal{G}$ embeds in a knot K if $\mathcal{G}$ can be completed by an arc exterior to S to form the knot K ; that is, there exists some arc in $S^3 - \text{Int}(S)$ such that upon gluing this arc to $\mathcal{G}$ along their boundary points, we have a knot in S^3 which is isotopic to K . We say that K is a closure of $\mathcal{G}$.

Let l denote a longitude for S which is contained in ∂S and avoids the genus-1 tangle. A closure K of $\mathcal{G}$ is called odd (respectively, even) with respect to l if $lk(K, l)$ is odd (respectively, even). If l is chosen to be the longitude which circles the central hole of S as in Fig. 1, and we span the longitude l by a disk Δ filling the hole, then $lk(K, l)$ is the number of transverse intersections counted with sign of the arc which completes $\mathcal{G}$ to K with Δ . Thus, in this case we can say more colloquially that K is an odd (respectively, even) closure with respect

Key words and phrases. Tangle, knot, branched cover, determinant.




FIGURE 1. Krebs' genus-1 tangle $\mathcal{A}$ in S together with a specified longitude l

to l if the arc which completes $\mathcal{G}$ to K passes through the hole of S an odd (respectively, even) number of times.

In [1], Krebs asks whether the genus-1 tangle given in Fig. 1 embeds in the unknot. We denote this tangle by $\mathcal{A}$, and when discussing this example, we always use the longitude l drawn in Fig. 1. Using the following results from [4], we are able to partially answer the question posed by Krebs.

Theorem 1.1 (Ruberman). *Suppose M is an orientable 3-manifold with connected boundary, and $i : M \hookrightarrow N$ where N is an orientable 3-manifold with $H_1(N)$ torsion. Then the inclusion map i_* induces an injection of the torsion subgroup $T_1(M)$ of $H_1(M)$ into $H_1(N)$.*

This theorem has a useful corollary which can easily be proved directly using a Meyer-Vietoris sequence.

Corollary 1.2 (Ruberman). *Let M and N be as in Theorem 1.1 but suppose $H_1(N) = 0$. Then $H_1(M)$ is torsion-free.*

One obtains an obstruction to embedding genus-1 tangles in knots from Theorem 1.1 by applying the result to branched covers of S branched over genus-1 tangles.

Recall, for a given n , each n -fold cover of S branched over a genus-1 tangle $\mathcal{G}$ is associated to a homomorphism $\varphi : H_1(S - \mathcal{G}) \rightarrow \mathbb{Z}_n$ which maps the meridian m of $\mathcal{G}$ to one. The remaining generator l of $H_1(S - \mathcal{G})$ may be sent to any element of $\mathbb{Z}_n$; we use $\varphi(l)$ to index the n -fold branched covers. So, $Y_{\mathcal{G},i}$ denotes the n -fold cover of S branched over $\mathcal{G}$ associated to the homomorphism φ which maps l to i .

If a genus-1 tangle $\mathcal{G}$ embeds in a knot K , then the n -fold cover X_K of S^3 branched over K restricts to some n -fold cover $Y_{\mathcal{G},i}$ of S branched over $\mathcal{G}$. In this case, we say that the closure K induces the cover $Y_{\mathcal{G},i}$. Then according to Theorem 1.1, the torsion subgroup $T_1(Y_{\mathcal{G},i})$ of $H_1(Y_{\mathcal{G},i})$ injects into $H_1(X_K)$.

Note that if K is the unknot, then X_K is S^3 and according to Corollary 1.2, the torsion subgroup $T_1(Y_{\mathcal{G},i})$ is trivial. Thus, if there is any




FIGURE 2. The disk in S where we perform a meridional twist, and the genus-1 tangle which results from the twist.

torsion in the homology of $Y_{\mathcal{G},i}$, then any closure of $\mathcal{G}$ which induces the cover $Y_{\mathcal{G},i}$ is not the unknot.

After applying this obstruction to the double-branched covers of S branched over $\mathcal{A}$, we prove the following results:

Theorem 1.3. *If a knot K in S^3 is an odd closure of $\mathcal{A}$, then $\det(K)$ is divisible by 3.*

Corollary 1.4. *If $\mathcal{A}$ embeds in the unknot, then the unknot is an even closure of $\mathcal{A}$.*

Before further discussion, we need to make a remark about the definition of genus-1 tangles.

Remark 1.5. Note that when defining genus-1 tangles, we fix a standardly embedded solid torus S in the 3-sphere. The reason that we restrict to a fixed embedding is that there are many ways to re-embed a solid torus inside S^3 .

For instance, if we perform a meridional twist on S along the disk indicated in Fig. 2, the image of $\mathcal{A}$ under this twist can be easily seen to embed in an unknot via the exterior arc pictured in Fig. 2. Thus it is necessary to specify the embedding of $S^1 \times D^2$ in the case of genus-1 tangles, and we restrict to a fixed standardly embedded solid torus in our definition.

2. SURGERY DESCRIPTIONS FOR DOUBLE-BRANCHED COVERS

For the purposes of this paper, we restrict our attention to double-branched covers of S branched over $\mathcal{A}$. Since a homomorphism $\varphi : H_1(S - \mathcal{A}) \rightarrow \mathbb{Z}_2$ must map the specified longitude l to either zero or one, there are two double-branched covers, $Y_{\mathcal{A},0}$ and $Y_{\mathcal{A},1}$. We call $Y_{\mathcal{A},0}$ the even double-branched cover because it is induced by all even




FIGURE 3. We perform surgery around a crossing, following [3].




FIGURE 4. Unwinding the genus-1 tangle $\mathcal{A}$ to make it look trivial. The surgery curve is always given the black-board framing.

closures of $\mathcal{A}$ (with respect to l). Similarly, since $Y_{\mathcal{A},1}$ is induced by all odd closures of $\mathcal{A}$, we call it the odd double-branched cover.

In this section, we adapt Rolfsen's technique to find surgery descriptions for these double-branched covers.

Following [3], we perform surgery near a carefully selected crossing (see Fig. 3) in such a way that after surgery we may essentially unwind $\mathcal{A}$ (via sliding its endpoints around the boundary in the complement of l) so that it looks trivial. This process, illustrated in Fig. 4, results in a nice surgery description of $\mathcal{A}$ inside S . Note that in the last drawing of Fig. 4, we choose to draw this surgery description in a particular way because it makes constructing branched covers easier.




FIGURE 5. Constructing the odd double-branched cover $Y_{\mathcal{A},1}$ of S branched over $\mathcal{A}$.



Now we construct the odd cover, $Y_{\mathcal{A},1}$. Construction is dictated by the homomorphism $\varphi : H_1(S - \mathcal{A}) \rightarrow \mathbb{Z}_2$ corresponding to the cover. If φ maps a generator of $H_1(S - \mathcal{A})$ to a non-zero element, then we cut the solid torus along a disk transverse to that generator. Thus, we have two cuts to make in the case of the odd cover.

First, we cut S along a disk which is transverse to the meridian m of $\mathcal{A}$ and whose boundary is made up of the unwound genus-1 tangle $\mathcal{A}$ together with an arc in ∂S . Then, because φ sends l to one, we cut S along a disk which is transverse to l and whose boundary is contained in ∂S . We then take two copies of the resulting manifold and glue them together carefully to obtain a surgery description for $Y_{\mathcal{A},1}$. This process is illustrated in Fig. 5.

Although it is not needed in the proof of Theorem 1.3, we also give a surgery description of the even double-branched cover $Y_{\mathcal{A},0}$ in Fig. 6.

3. HOMOLOGY OF THE COVERS

Now we compute the homology of the odd double-branched cover. From Fig. 5 we see that the surgery description for $Y_{\mathcal{A},1}$ is given by

FIGURE 6. Obtaining a surgery description of $Y_{A,0}$.FIGURE 7. A surgery description of $Y_{A,1}$

a 2-component surgery link inside a genus-2 handlebody. We denote the components of the surgery link by σ and τ , and let H denote the genus-2 handlebody. The complement of H in S^3 is a neighborhood of the handcuff graph G , pictured in Fig. 7, which is composed of loops α_1 and α_2 joined together by an arc. Then the complement of $\sigma \cup \tau$ in H can be viewed as the complement of $\sigma \cup \tau \cup G$ in S^3 . One can see that $H_1(S^3 - (\sigma \cup \tau \cup G))$ is isomorphic to $H_1(S^3 - (\sigma \cup \tau \cup \alpha_1 \cup \alpha_2))$ which is free on four generators: the meridians of σ , τ , α_1 , and α_2 .

Completing the surgery by gluing in two solid tori according to σ and τ introduces two relations on these four generators, which are given by the linking numbers of σ and τ with each of σ , τ , α_1 , and α_2 . Then $H_1(Y_{A,1})$ is isomorphic to $H_1(S^3 - (\sigma \cup \tau \cup \alpha_1 \cup \alpha_2))$ modulo these two relations, and we can get a presentation for $H_1(Y_{A,1})$ using linking

numbers. Thus, we have the following presentation matrix for $H_1(Y_{\mathcal{A},1})$:

$$\begin{array}{c} \sigma \quad \tau \quad \alpha_1 \quad \alpha_2 \\ \sigma \begin{pmatrix} 1 & 2 & 0 & 0 \\ 2 & 1 & 0 & 0 \end{pmatrix} \\ \tau \end{array}$$

Using row and columns operations we obtain a simpler presentation matrix:

$$\begin{array}{c} \sigma \quad \tau \quad \alpha_1 \quad \alpha_2 \\ \sigma \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \end{pmatrix} \\ \tau \end{array}$$

Therefore, $H_1(Y_{\mathcal{A},1}) = \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}_3$ and we are now able to prove the main theorem. Corollary 1.4 follows immediately.

Proof of Theorem 1.3. Let K be an odd closure of $\mathcal{A}$, and let X_K denote the double cover of S^3 branched over K . Since K is an odd closure of $\mathcal{A}$, it induces a restriction from X_K to $Y_{\mathcal{A},1}$. Then according to Theorem 1.1, we have that $T_1(Y_{\mathcal{A},1}) = \mathbb{Z}_3 \hookrightarrow H_1(X_K)$. Thus $|T_1(Y_{\mathcal{A},1})| = 3$ divides $|H_1(X_K)| = \det(K)$. $\square$

We are unable to use this method to restrict all closures of $\mathcal{A}$ because $Y_{\mathcal{A},0}$ has a torsion-free first homology group. Indeed, the statement in Remark 1.5 allows us to see that the even cover does embed in S^3 and so must have torsion-free first homology. Of course, this can be verified by deriving a presentation for the homology of $Y_{\mathcal{A},0}$ using the procedure above.

REFERENCES

- [1] D. A. Krebs, An obstruction to embedding 4-tangles in links, *J. Knot Theory and its Ramifications* **8** (1999) 321-352.
- [2] J.H. Przytycki, D.S. Silver and S.G. Williams, 3-manifolds, tangles, and persistent invariants, *Math. Proc. Camb. Phil. Soc.* **139** (2005) 291-306.
- [3] D. Rolfsen, *Knot and Links* (Publish or Perish, Berkeley, CA, 1976).
- [4] D. Ruberman, Embedding tangles in links, *J. Knot Theory and its Ramifications* **9** (2000) 523-530.

MATHEMATICS DEPARTMENT, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA

E-mail address: `sabern1@tigers.lsu.edu`